



**NAMIBIA UNIVERSITY
OF SCIENCE AND TECHNOLOGY
FACULTY OF HEALTH AND APPLIED SCIENCES
DEPARTMENT OF MATHEMATICS AND STATISTICS**

QUALIFICATION: HEALTH INFORMATION SYSTEMS MANAGEMENT	
QUALIFICATION CODE: 07BHIS	LEVEL: 7
COURSE CODE: BSD721S	COURSE: Biostatistics and Demography
SESSION: January 2019	PAPER : THEORY
DURATION: 3 Hours	MARKS: 100

SECOND OPPORTUNITY EXAMINATION QUESTION PAPER	
EXAMINER	Mr. J. J. SWARTZ
MODERATOR:	Dr. L. PAZVAKAWAMBWA

INSTRUCTIONS
<ol style="list-style-type: none">1. Answer all the questions in the booklet provided2. Show clearly all the steps used in the calculations.

PERMISSIBLE MATERIALS

Calculator

APPENDICES

Statistical Tables

THIS QUESTION PAPER CONSISTS OF 4 PAGES (Including this front page)

Question 1 [25]

1.1. Choose the right answer:

1.1.1. Which of the following is not an example of discrete variable? [2]

- A. the number of students at the class of statistics.
- B. the number of times a child cry in a certain street.
- C. the time to run a certain distance.
- D. the number of buildings in a certain street.
- E. number of educated persons in a family.

1.1.2. Which of the following is an example of qualitative variable? [2]

- A. the blood pressure level.
- B. the number of times a child brush his/her teeth.
- C. whether or not someone fail in an exam.
- D. Weight of babies at birth.
- E. the time to run a certain distance

1.2. The following table shows the number of hours 45 hospital patients slept following the administration of a certain anaesthetic.

Class Interval	Frequency
1-5	21
6-10	16
11-15	6
16-20	2
Total	45

From these data construct:

1.2.1. A relative frequency distribution. [2]

1.2.2. How many of the measurements are greater than 10? [1]

1.2.3. What percentage of the measurements are between 6-15? [1]

1.2.4. What proportion of the measurement is less than or equal 15? [1]

1.3. Cardosi et al. (A-12) performed a 4 years retrospective review of 102 women undergoing radical hysterectomy for cervical or endometrial cancer. Catheter-associated urinary tract infection was observed in 12 of the subjects. Below are the numbers of postoperative days until diagnosis of the infection for each subject experiencing an infection.

16	10	49	15	6	15	8	19	11	22	13	17
----	----	----	----	---	----	---	----	----	----	----	----

Use the data above and find the

1.4.1 Mean [2]

1.4.2 Median [2]

1.4.3 Mode [1]

1.4.4 Range [2]

1.4.5 Variance [3]

1.4.6 Standard deviation. [2]

1.4.7 Coefficient of variation. [2]

Question 2 [25]

2.1. Provide four non-probability sampling methods [4]

2.2. The Namibia Statistics Agency (NSA) wants to estimate the rate of poverty among the households in Namibia. How large a sample should be taken to be 95% confident that the error of estimation of the poverty levels does not exceed 0.05? The true value of p is expected to be near 0.17. [6]

2.3. A pharmacy store manager MoHSS is concerned that sales are being lost due to stock outs while waiting for new stock of morphine liquid. It has been determined that demand while waiting for new stock is normally distributed with a mean of 15 litres and a standard deviation of 6 litres.

2.3.1. The manager would like to know the probability of the demand below 20 litres during stockout. [5]

2.3.2. The manager would like to know the probability of the demand exceeding 20 litres during stockout. [5]

2.3.2. If the manager of wants the probability of demand during stock out to be no more than 0.05, what should the reorder point be? [5]

Question 3 [25]

3.1. Define the following terms:

3.1.1. Estimator [2]

3.1.2. Estimate [2]

3.2. The activity values of a certain enzyme measured in normal gastric tissue of 40 patients with gastric carcinoma has a mean of 0.718 and a standard deviation of 0.511. Construct a 90 % confidence interval for the population mean. [5]

3.3. NHP collected data on a sample of 301 women living in Windhoek. One variable of interest was the percentage of subjects with impaired fasting glucose (IFG). In the study, 24 women were classified in the (IFG) stage. The article cites population estimates for (IFG) among women in Windhoek as 6.3 percent. Is there sufficient evidence to indicate that the population of women in Windhoek has a prevalence of IFG higher than 6.3 percent. [10]

3.4. The following table below provide data on chest circumference (cm) and birthweight (kg) of 10 babies.

X(cm)	y(kg)	x ²	y ²	xy
22.4	2.00	501.76	4.00	44.8
27.5	2.25	756.25	5.06	61.88
28.5	2.10	812.25	4.41	59.85
28.5	2.35	812.25	5.52	66.98
29.4	2.45	864.36	6.00	72.03
29.4	2.50	864.36	6.25	73.5
30.5	2.80	930.25	7.84	85.4
32.0	2.80	1024.0	7.84	89.6
31.4	2.55	985.96	6.50	80.07
32.5	3.00	1056.25	9.00	97.5
TOTAL				
292.1	24.8	8607.69	62.42	731.61

3.4.1. Calculate the correlation coefficient between the chest circumference (cm) and birthweight (kg) and interpret your results. [6]

Question 4 [25]

4.1. Provide three basic components of population change. [3]

4.2 The table below presents Number of births to women of Namibia in urban and rural areas in the 12 months before the census, obtained from the 2001 Population and Housing Census of Namibia (Source: CBS, 2001 Population and Housing Census).

Age of Mother	Number of Women			Number of Births		
	Urban	Rural	Total	Urban	Rural	Total
15 - 19	30482	72509	102991	1277	3901	5278
20 - 24	36109	51993	88102	3827	8137	11964
25 - 29	36319	40160	76479	4389	6667	11056
30 - 34	28461	32943	61404	3217	5212	8429
35 - 39	22550	28783	51333	1793	3499	5292
40 - 44	16186	23694	39880	613	1770	2383
45 - 49	10961	19846	30807	120	561	681
Total	181068	269928	450996	15336	29747	45083

4.2.1) Calculate the Crude Birth Rate (CBR) for the total population of Namibia. Use the midyear population, P = 1 830 330 [3]

4.2.2) Calculate the General Fertility Rate (GFR) [3]

4.2.3) Calculate the age-specific fertility (ASFR) rates for each age cohort [7]

4.2.4) Calculate the Total Fertility Rate (TFR) for the total population of Namibia [4]

4.2.5) Calculate the Child Women Ratio (CWR) for the total population of Namibia. Use the midyear population, P = 1 830 330 with child (0 – 4) years population of 98 460. [3]

4.3 Briefly define a life table, and discuss the importance of a life table [2]

*****END OF EXAMINATION*****

Standard Normal Probabilities




Table entry for z is the area under the standard normal curve to the left of z .

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

t Table

cum. prob	<i>t</i> _{.50}	<i>t</i> _{.75}	<i>t</i> _{.80}	<i>t</i> _{.85}	<i>t</i> _{.90}	<i>t</i> _{.95}	<i>t</i> _{.975}	<i>t</i> _{.99}	<i>t</i> _{.995}	<i>t</i> _{.999}	<i>t</i> _{.9995}
one-tail	0.50	0.25	0.20	0.15	0.10	0.05	0.025	0.01	0.005	0.001	0.0005
two-tails	1.00	0.50	0.40	0.30	0.20	0.10	0.05	0.02	0.01	0.002	0.001
df											
1	0.000	1.000	1.376	1.963	3.078	6.314	12.71	31.82	63.66	318.31	636.62
2	0.000	0.816	1.061	1.386	1.886	2.920	4.303	6.965	9.925	22.327	31.599
3	0.000	0.765	0.978	1.250	1.638	2.353	3.182	4.541	5.841	10.215	12.924
4	0.000	0.741	0.941	1.190	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	0.000	0.727	0.920	1.156	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	0.000	0.718	0.906	1.134	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	0.000	0.711	0.896	1.119	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	0.000	0.706	0.889	1.108	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	0.000	0.703	0.883	1.100	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	0.000	0.700	0.879	1.093	1.372	1.812	2.228	2.764	3.169	4.144	4.587
11	0.000	0.697	0.876	1.088	1.363	1.796	2.201	2.718	3.106	4.025	4.437
12	0.000	0.695	0.873	1.083	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	0.000	0.694	0.870	1.079	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	0.000	0.692	0.868	1.076	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	0.000	0.691	0.866	1.074	1.341	1.753	2.131	2.602	2.947	3.733	4.073
16	0.000	0.690	0.865	1.071	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	0.000	0.689	0.863	1.069	1.333	1.740	2.110	2.567	2.898	3.646	3.965
18	0.000	0.688	0.862	1.067	1.330	1.734	2.101	2.552	2.878	3.610	3.922
19	0.000	0.688	0.861	1.066	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	0.000	0.687	0.860	1.064	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	0.000	0.686	0.859	1.063	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	0.000	0.686	0.858	1.061	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	0.000	0.685	0.858	1.060	1.319	1.714	2.069	2.500	2.807	3.485	3.768
24	0.000	0.685	0.857	1.059	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	0.000	0.684	0.856	1.058	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	0.000	0.684	0.856	1.058	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	0.000	0.684	0.855	1.057	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	0.000	0.683	0.855	1.056	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	0.000	0.683	0.854	1.055	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	0.000	0.683	0.854	1.055	1.310	1.697	2.042	2.457	2.750	3.385	3.646
40	0.000	0.681	0.851	1.050	1.303	1.684	2.021	2.423	2.704	3.307	3.551
60	0.000	0.679	0.848	1.045	1.296	1.671	2.000	2.390	2.660	3.232	3.460
80	0.000	0.678	0.846	1.043	1.292	1.664	1.990	2.374	2.639	3.195	3.416
100	0.000	0.677	0.845	1.042	1.290	1.660	1.984	2.364	2.626	3.174	3.390
1000	0.000	0.675	0.842	1.037	1.282	1.646	1.962	2.330	2.581	3.098	3.300
Z	0.000	0.674	0.842	1.036	1.282	1.645	1.960	2.326	2.576	3.090	3.291
	0%	50%	60%	70%	80%	90%	95%	98%	99%	99.8%	99.9%
	Confidence Level										