

NAMIBIA UNIVERSITY

OF SCIENCE AND TECHNOLOGY

FACULTY OF HEALTH, APPLIED SCIENCES AND NATURAL RESOURCES

DEPARTMENT OF NATURAL AND APPLIED SCIENCES

QUALIFICATION: BACHELOR OF SCIENCE	CE HONOURS
QUALIFICATION CODE: 08BOSH	LEVEL: 8
COURSE CODE: BBC811S	COURSE NAME: BIOINORGANIC AND BIOPHYSICAL CHEMISTRY
SESSION: JULY 2022	PAPER: THEORY
DURATION: 3 HOURS	MARKS: 100

SUPPLEME	NTARY/SECOND OPPORTUNITY EXAMINATION QUESTION PAPER
EXAMINER(S)	DR. EUODIA HESS
MODERATOR:	DR. LIKIUS DANIEL

	INSTRUCTIONS
1.	Answer ALL the questions.
2.	Write clearly and neatly.
3.	Number the answers clearly
4.	All written work must be done in blue or black ink and sketches can
	be done in pencil
5.	No books, notes and other additional aids are allowed

THIS QUESTION PAPER CONSISTS OF 4 PAGES (Including this front page, list of useful constants and Periodic Table)

QUESTION 1:	[10]
What are the biological roles/functions of the following metals: a) Co b) Cu c) Na d) Ca e) K	
QUESTION 2:	[10]
What are the donor atoms according to the hard soft acid base (HSAB) theory? Explain with examples if they hard, soft or intermediate.	
QUESTION 3:	[40]
3.1 In medicine the use of metal ions and their associated complexes is widespread. Among Metal ions commonly used over the centuries were Hg ²⁺ for treatment of syphilis, Mg ²⁺ for intestinal disorders and Fe ²⁺ for anaemia. Today one of the leading anticancer drug is cis-[Pt(NH ₃) ₂ Cl ₂], cisplatin, which was first approved for use in 1978. i) Against which type(s) of cancer is cisplatin effective? ii) Draw the structure of cisplatin. iii) How does cisplatin function as a anticancer drug? iv) What are the disadvantages of using cisplatin as an anticancer drug? 3.2 Name three other platinum drugs that are utilized as an anticancer drug and for each drug answer question i – iv for them as well.	(3) (2) (3) (2) (30)
SECTION B:	[40]

The kinetics of CO₂ hydration catalysed by the enzyme carbonic anhydrase is as follows:

 $CO_2 + H_2O \leftrightarrow HCO_3^- + H^+$

The following initial reaction rates for the hydration reaction were obtained for an initial enzyme concentration of 2.3 nM and temperature of 0.5 $^{\circ}$ C:

Rate (M s-1)	[CO ₂] (mM)	
2.78 x 10 ⁻⁵	1.25	
5.00 x 10 ⁻⁵	2.5	
8.33 x 10 ⁻⁵	5.0	
1.67 x 10 ⁻⁴	20.0	

QUESTION 2: [10]

2.1 Obtain a balanced chemical equation and calculate the standard electrochemical potential for reduction of Fe³⁺ to Fe by Zn metal (Zn²⁺ \rightarrow Zn E⁰= - 0.7618 V; Fe³⁺ \rightarrow Fe E⁰ = - 0.037 V). (5)

2.2 The enzyme glutathione reductase replenishes the cell's supply of GSH regenerating two molecules of GSH from single molecule of oxidized gluthione (GSSH), using NADPH as a source of two reducing equivalents. A typical cellular NADP+/NADPH ratio is 0.005, calculate the equilibrium cellular concentration of GSSG at pH 7 and 25°C, if GSH concentration is 4 mM. (GSSG + 2H+ + 2e- → 2 GSH E0 = -0.240 V; NADP+ + H+ + 2e- → NADPH E0 = -0.339 V). (5)

QUESTION 3: [10]

For the hydrolysis of ATP, standard conditions do not prevail in the cellular environment. In the Cell, typical concentrations of ATP, ADP and inorganic phosphate are $c_{\text{ATP}} = 1850 \ \mu\text{M}$, $c_{\text{ADP}} = 138 \ \mu\text{M}$, and $c_{\text{pi}} = 1.00 \ \text{mM}$. Calculate the Gibbs energy of hydrolysis in the cellular environment, assuming pH = 7 and T = 310 K.

QUESTION 4: [10]

The enthalpy of melting ice at 1 bar is 6.007 kJ/mol; the density of water at 0 °C is 999.9 kg m⁻³, While that of ice is 915.0 kg m⁻³. Assuming ΔV_m^{fusion} and ΔH_m^{fusion} are constant, determine the freezing point of water at 100 bar.

END OF EXAMINATION

USEFUL CONSTANTS:

Gas constant, $R = 8.3145 \text{ J} \cdot \text{mol}^{-1} \cdot \text{K}^{-1} = 0.083145 \text{ dm}^3 \cdot \text{bar} \cdot \text{mol}^{-1} \cdot \text{K}^{-1} = 0.08206 \text{ L atm mol}^{-1} \cdot \text{K}^{-1} = 0.08206 \text{ L}$

 $1 \text{ Pa} \cdot \text{m}^3 = 1 \text{ kPa.L} = 1 \text{ N} \cdot \text{m} = 1 \text{ J}$

1 atm = 101 325 Pa = 760 mmHg = 760 torr

Faradays constant = 96,485 C/mol

Avogadro's Number, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

Planck's constant, $h = 6.626 \times 10^{-34} \text{ Js}$

Speed of light, $c = 2.998 \times 10^8 \text{ ms}^{-1}$

PERIODIC TABLE OF THE ELEMENTS

1		1		_				_			_	-	_					_	_
24 Be 9.01218 Ca Sc Ti V Ca Sc Ti Nb H H T Ta H T T	18 2 He	10		20.179	18	Ar	39.948	36	Kr	83.8	54	Xe	131.29	98	Rn	(222)	118	Uuo	
24 Be 9.01218 Rg 4	17	6	H	18.9984	17	Ü	35.453	35	Br	79.904	53	_	126.9	85	At	(210)			
24 Be 9.01218 12 12 13 14 14 18 18 19 19 19 19 19 19 19 19 19 19 19 19 19	16	8	0	15.9994	16	Ø	32.06	34	Se		52	Te	127.6	84	Po	(509)	116	Unh	
24 Be 9.01218 12 12 13 14 14 18 18 19 19 19 19 19 19 19 19 19 19 19 19 19	15	7	Z	14.0067	15	Д	30.9738	33	As	74.9216	51	Sb	121.75	83	Bi	208.908			
2 Be 9.01218 12 3 4 5 6 7 8 9 10 11 12 24.305 3 4 5 6 7 8 9 10 11 12 20 21 22 23 24 25 26 27 28 29 30 40.08 44.9559 47.88 50.9415 51.996 54.9380 55.847 58.9332 58.69 65.346 65.38 Sr Y Cr Mn Fe Co Ni Cr Zn 8.6 7 8 9 10 11 12 8.7 1 2 23 24 25 26 37 88 8.7 Y Cr Mn Fe Co Ni Pd 48 8.6 7 8 4 4 45 46 47 48	14	9	Ü	12.011	14	Si		32	g		50	Sn	118.69	82	Pb	207.2	114	Unq	
2 Mg 24.305 3 4 5.01218 La	13	S	В	10.81	13	ΙV	26.9815	31	g _B	69.72	64	In	114.82	18	E	204.383			
2 Mg 24.305 3 4 5.01218 La							12	30	Zn	65.38					Hg	200.59	112	Uub	(269)
2 Mg 24.305 3 4 5.01218 La							11	29	Cn	63.546	47	Ag	107.868	79	Au	196.961	111	Unn	(272)
2 4 Be 9.01218 Ag 24.305 3 4 5 7 7 8 7 8 7 8 7 8 7 8 8 8 8 103 174.967 178.49 180.948 183.85 186.207 190.2 8 8 103 104 105 105 106 107 108 108 108 108 108 108 108 108 108 108							10	28	Z	58.69		Pd	106.4				110	Unn	(269)
2 4 Be 9.01218 Ag 24.305 3 4 5 6 7 20 21 22 23 24 25 Ca Sc Ti V Cr Mn 40.08 44.9559 47.88 50.9415 51.996 54.9380 38 39 40 41 42 43 Sr Y Zr Nb Mo Tc 87.62 88.9059 91.22 92.9064 95.94 (98) 56 71 72 73 74 75 Ba Lu Hf Ta W Re 137.33 174.967 178.49 180.948 183.85 186.207 88 103 104 105 106 107 Ra Lr Rf Db Sg Bh 5.026.025 (260) (261) (262) (263) (264) (6	27	ථ	58.9332	45	Rh	102.906	11	Ir	192.22	109	Mt	(268)
2 4 Be 9.01218 Ag 24.305 3 4 5 Ca Sc Ca							∞	26	Fe	55.847	44	Ru	101.07	9/	SO	190.2	108	Hs	(265)
2 4 Be 9.01218 AMg 24.305 3 4 524.305 3 4 4 526.025 3 4 4 5 7 7 7 7 7 7 7 7 7 7 7 8 8 8 103 104 105 105 105 105 105 105 105 105 105 105							7	25	Mn	54.9380	43	Lc	(86)	75	Re	186.207	107	Bh	
2 4 Be 9.01218 24.305 3 4 22.305 3 4 20 21 22 Ca Sc Ti 40.08 44.9559 470 Sr Y Sr N Sr N Sr N Sr N Sr						,	9	24	Ċ		42		_	74	≩	183.85	106	S	(263)
2 4 Be 9.01218 12 Mg 24.305 3 4 20 21 22 Ca Sc Ca Sc Ti 88 103 104 Sc Ti 173.33 174.967 178.44 Ra Lu Hf Ra Lr Rd Rd Lc Rd Rd Lc Rd						,	2	23	>	50.9415	41	Sp	92.9064	73	Ta	180.948	105	Dp	(262)
2 4 4 Be 9.01218 12 12 20 Ca 40.08 38 88 88 Ra 226.025 226.025 26 26 26 26 26 26 26 26 26 26 26 26 26						,	4	22	Ë	_	40		91.22	72	HL	4	104	Rf	(261)
						,	3	21	Sc	44.9559	39	X	88.9059	71	Lu		103	Ľ	(260)
1 H 1.00794 3 Li 6.941 11 Na 22.9898 19 K 37 Rb 85.4678 55 Cs 132.905 87 Fr (223)	7	4	Be	9.01218	12	Mg	24.305	20	రొ	1	38	Sr		99	Ba	137.33	88	Ra	226.025
	1 H 1.00794	3	<u>5</u>	6.941	=	Na	22.9898	19	×	39.0983	37	Rb	85.4678	55	S	132.905	87	Fr	(223)

Londhonidon	6.7	02	02	3		3	"	1	1	1	-	1		
Lanuaniues;	<u>'</u>	28	29	00		79	63	49	65	99	67	89	69	70
	La	ပီ	Pr	PN	Pm	Sm	Sm Eu	S	Tb	Dy	Dy Ho Er	Er	Lm	Yb
	138.906	140.12	140.12 140.908 1	144.24		150.36	151.96	57.2	58.92	162.50	161.930	167.26	6.934	173.04
Actinides:	68	06	91	92	93	94	95	96	26	86	66	100	101	102
	Ac	Th	Pa	D	Sp	Pu	Am	Cm	Bk	C	Es	Fm	Md	S
	227.028	232.038	227.028 232.038 231.036 238.029 237.048	238.029	237.048	(244)	(243)	(247)	(247)	(251)	(252)	(257)		(259)