

FACULTY OF ENGINEERING

InSTEM

QUALIFICATION: INTRODUCTION TO SCIENCE, TECHNOLOGY, ENGINEERING AND MATHEMATICS					
QUALIFICATION CODE: 04STEM		LEVEL: 4			
COURSE CODE: IPH401S		COURSE NAM	COURSE NAME: INTRODUCTION TO PHYSICS A		
SESSION:	JANUARY 2020	PAPER:	N/A		
DURATION:	3 HOURS	MARKS:	100		

S	ECOND OPPORTUNITY EXAMINATION QUESTION PAPER		
EXAMINER(S) Ms Ilana Malan			
	Ms Oksana Kachepa		
MODERATOR:	Mr Anthony Apata		

INSTRUCTIONS

- 1. Answer all questions.
- 2. Write all the answers in ink.
- 3. No books, notes, correction fluid (Tippex) or cell phones allowed.
- 4. Pocket calculators are allowed.
- 5. You are not allowed to borrow or lend any equipment or stationary.
- 6. All FINAL ANSWERS must be rounded off to THREE DECIMAL PLACES.
- 7. All CONSTANT VALUES and FORMULAS on page 11.
- 8. Use GRAPH PAPER provided for question 16.4

THIS QUESTION PAPER CONSISTS OF 11 PAGES (Excluding this front page)

SECTION A - TOTAL MARKS 30

This section consists of nine questions. Choose the correct answer and clearly indicate your answer on your answer sheet.

Question 1 [4]

How much work is required to accelerate a 2.5 t car from 5 km/h to 60 km/h?

- (a) 291.763 kJ
- (b) 344.81 kJ
- (c) 3781.25 kJ
- (d) 347.222 kJ

Question 2 [4]

In which quadrant will the equilibrant force of the following be:

$$\Sigma_{x} = -3 + 2\cos 10^{\circ}$$
 and $\Sigma_{y} = -3 + 2\cos 10^{\circ}$

- (a) first
- (b) second
- (c) third
- (d) fourth

Question 3 [4]

A 410 g non-elastic ball rolling at 5.2 m/s collides with a brick wall and comes to a standstill in 41 ms. Calculate the force required.

- (a) 126.829 N
- (b) 0.874 N
- (c) 52 N
- (d) 309.3396 N

Question 4 [2]

Choose the correct expression for the First Law of Thermodynamics:

- (a) $\Delta U = \Delta Q + \Delta W$
- (b) $\Delta Q = \Delta U + \Delta W$
- (c) $\Delta W = \Delta Q + \Delta U$
- (d) $\Delta Q \Delta U = \Delta W$


Question 5 [2]

Which of the following is NOT a vector quantity?

- (a) kinetic energy
- (b) pressure
- (c) acceleration
- (d) momentum

Question 6 [2]

Four objects are standing on a flat surface in **Figure 1**. M indicates the centre of mass. Which object will fall over?



- (a) A
- (b) B
- (c) C
- (d) D

Question 7 [4]

Figure 3 shows the velocity-time graph for a moving object. Determine the total distance moved from t = 4 s to t = 13 s.




Figure 2

- (a) 12 m
- (b) 99 m
- (c) 87 m
- (d) 111 m

Question 8 [4]

A large heavy steel ball is dropped from a height of 75 m. At the same time when the ball is released the timer starts. How far does the ball from t = 0.5 s to t = 2.7 s?

- (a) 34.5312 m
- (b) 75 m
- (c) 35 m
- (d) 23.7402 m

Question 9

5

[4]

Determine the phase difference in degrees between the two waves in Figure 3.




Figure 3

- (a) 23°
- (b) 43°
- (c) 63°
- (d) 83°

SECTION B - TOTAL MARKS 70

This section consists of six (6) questions. Answer ALL the questions.

Question 10 [10]

Answer the questions 10.1 - 10.4 with regards to **Figure 4**.




Figure 4

10.1	When is the object stationary?	(1)
10.2	When is the object accelerating?	(1)
10.3	Describe the movement from $t = 0$ s to $t = 6$ seconds.	(3)
10.4	Determine total distance.	(2.5)
10.5	Determine total displacement.	(2.5)

Question 11 [10]

The 17 m long flagpole with a mass of 35 kg, as shown in **Figure 5**, has a non-uniform shape. It is attached to a 10 m vertical pole on the wider side of the flagpole. The centre of mass of the flagpole is a third of the length of the flagpole from the left side. A flag with a mass of 14 kg hangs vertical from the right end. A strong steel cable is attached at the centre of the flagpole at an angle of 46° to the horizontal.




Figure 5

- 11.1 Determine the tension, T, in the cable for the flagpole to be in horizontal equilibrium. (6)
- 11.2 If the flag is moved to 10 m from the left end, what will the tension in the cable be? (4)

Question 12	[12]

Two projectiles are launched at the same time at different angles and different initial velocities. The reach the same maximum height. Projectile A is released at an angle of 34° to the horizontal with an initial velocity of 35 m/s.

12.1	Projectile B is launched at an initial velocity of 45 m/s. Determine the angle with	
	the horizontal at which projectile B is launched.	(3)
12.2	Determine the difference in range when the two projectiles strike the ground.	(6)
12.3	How high is projectile A after 3 seconds?	(3

Question 13 [10]

A train carrying passengers to a ski resort in the Alps has a total mass of 55.60 t. The train maintains a constant speed of 28 km/h. The track is horizontal for 42 km and then has a steep incline towards the resort for another 5 km. The speed for the last 5 km is constant at 3 km/h.

13.1	Determine the work done over the first half of the journey.	(2
13.2	The work done over the last 5 km of the journey is 2 392. 333 MJ with the train	
	experiencing a frictional force of 4 N/kg. Determine the angle of the incline.	(5)
13.3	How long will the journey take?	(3)

Question 14 [8]

- 14.1 Give one word for: "Turning effect of a force." (1)
- 14.2 The lever in **Figure 6** of length 25 m, is in a horizontal equilibrium. The centre of gravity is in the middle of the lever. Determine the angle B. (7)




Figure 6

Question 15 [12]

- 15.1 Describe latent heat of a substances. (2)
- 15.2 The graph in **Figure 7** compares the energy in a substance as the temperature decreases. This graph is for an unknown substance of mass 40 g with a boiling point of 150°C. Heat is given out at a constant rate of 15 W.




Figure 7

15.2.1 How much heat was released during condensation? (1)
15.2.2 Determine the latent heat of vaporization. (2)
15.2.3 Determine the specific heat capacity_{liquid} of the substance. (2)
15.2.4 How long will it take cool down 2 kg of the substance that has just condensed, to freezing point? (3)

Question 16 [8]

Answer all the questions with reference to Figure 8.

The object makes just ONE oscillation according to the sequence: A - B - A - C - A.




Figure 8

What is the direction of the restoring force at C? (1)
At which position(s) will the object have maximum potential energy? (1)
What sign will the velocity have at point A and B, while in movement from A to B? (1)
Draw a graph showing how the kinetic energy varies for one oscillation. (5)

The End

Constants and formulas:

average speed =
$$\frac{\text{distance}}{\text{time}}$$

average velocity =
$$\frac{u+v}{2}$$

$$s = ut + \frac{1}{2}at^2$$

$$F = ma$$

$$E_p = mgh$$

$$\eta = \frac{\text{output}}{\text{input}} \times 100\%$$

 $power = F \times velocity$

$$F = \frac{\Delta momentum}{\Delta t} = \frac{\Delta p}{\Delta t}$$

$$\Delta U = \Delta Q + \Delta work$$

$$E = mL$$

$$\frac{p_1 V_1}{T_1} = \frac{p_2 V_2}{T_2}$$

10 ³	kilo	K	10-2	centi	С
10 ⁶	Mega	М	10-3	milli	m
10 ⁹	Giga	G	10-6	micro	μ
10 ¹²	Tera	Т	10 ⁻⁹	nano	n
10 ¹⁵	Peta	Р	10-12	pico	р

$$g = 9.81 \text{ m/s}^2$$
 0 K = -273,15 °C

$$0 \text{ K} = -273,15 \, ^{\circ}\text{C}$$

average velocity =
$$\frac{\text{displacement}}{\text{time}}$$

$$a = \frac{v - u}{t}$$

$$v^2 = u^2 + 2as$$

$$moment = Fs$$

pressure =
$$\frac{F}{A}$$

$$E_k = \frac{1}{2}mv^2$$

$$power = \frac{work}{t}$$

$$m_1 u_1 + m_2 u_2 = m_1 v_1 + m_2 v_2$$

$$f = \frac{1}{T}$$

$$E/Q = mc\Delta T$$

$$pV = nRT$$

work =
$$pAs = p\Delta V$$