



**NAMIBIA UNIVERSITY
OF SCIENCE AND TECHNOLOGY**

**Faculty of Health, Natural
Resources and Applied
Sciences**

**School of Natural and Applied
Sciences**

**Department of Mathematics,
Statistics and Actuarial Science**

13 Jackson Kaujeva Street
Private Bag 13388
Windhoek
NAMIBIA

T: +264 61 207 2913
E: msas@nust.na
W: www.nust.na

QUALIFICATION : BACHELOR OF SCIENCE HONOURS IN APPLIED STATISTICS	
QUALIFICATION CODE: 08BSHS	LEVEL: 8
COURSE: SAMPLING THEORY	COURSE CODE: SAT802S
DATE: NOVEMBER 2024	SESSION: 1
DURATION: 3 HOURS	MARKS: 100

FIRST OPPORTUNITY: QUESTION PAPER

EXAMINER: *Mr. Jan Johannes Swartz*

MODERATOR: *Prof. Opeoluwa Oyedele*

INSTRUCTIONS:

1. Answer all questions on the separate answer sheet.
2. Please write neatly and legibly.
3. Do not use the left side margin of the exam paper. This must be allowed for the examiner.
4. No books, notes and other additional aids are allowed.
5. Mark all answers clearly with their respective question numbers.

PERMISSIBLE MATERIALS:

1. Non-Programmable Calculator

ATTACHMENTS

1. Z - Table
2. T - Table

This paper consists of 5 pages including this front page

Question 1 [25 marks]

1.1 Write a short description on the importance of the normal distribution in sampling theory. [4]

1.2 Provide six basic steps in developing a sampling plan. [6]

1.3 For the 200 managers and 800 engineers of a corporation, the standard deviations of the number of days a year spent on research were presumed to be 30 and 60 days, respectively. Find the sample size needed for proportional allocation to estimate the population mean with the standard error of the estimator not exceeding 10 and its allocation for the two groups. [5]

1.4 Among 100 Retailers in Namibia, the average of employee sizes for the largest 10 and smallest 10 corporations were known to be 300 and 100, respectively. For a sample of 20 from the remaining 80 retailers, the mean and standard deviation were 250 and 110, respectively. For the total employee size of the 80 retailers, find the

1.4.1 Estimate for the total, [2]

1.4.2 Standard Error of the estimate, and [3]

1.4.3 95% confidence limits. [5]

Question 2 [25 marks]

2.1. The Ministry of Health and Social Services (MoHSS) wants to estimate the rate of incidence of respiratory disorders among the middle-aged male and female smokers in Namibia. How large a sample should be taken to be 95% confident that the error of estimation of the proportion of the population with such disorders does not exceed 0.05? The true value of p is expected to be near 0.30. [5]

2.2.

We propose to estimate the mean \bar{Y} of a characteristic y by way of a sample selected according to a simple random design without replacement of size 1000 in a population of size 1 000 000. We know the mean $\bar{X} = 1.5$ of an auxiliary characteristic x . We have the following results:

$$s_y^2 = 2.0, s_x^2 = 2.5, s_{xy} = 1.5, \hat{X} = 1.4, \hat{Y} = 1.0$$

2.2.1. Estimate \bar{Y} by way of Horvitz – Thompson, difference, ratio and regression estimators. Estimate the variances of these estimators. [15]

2.2.2. Which estimator should we choose to estimate \bar{Y} ? [5]

Question 3 [25 marks]

3.1. The Namibian 25, 2001, summarized the results of a survey conducted by Yellow Express on 2000 lawyers on sexual advances in the office. Between 85 and 98% responded to the questions in the survey; 49% of the responding women and 9% of the responding men agreed that some sorts of harassment exist in the offices. Assume that the population of lawyers is large and there are equal numbers of female and male lawyers, and ignore the nonresponse; that is, consider the respondents to be a random sample of the 2000 lawyers.

3.1.1 Find the standard errors for females and males. [5]

3.2. A forest resource manager is interested in estimating the total number of dead trees in a 400-acre area of heavy infestation. She subdivides the area into 200 plots of equal sizes and uses photo counts to find the number of dead trees in 18 randomly sampled plots. She then randomly samples 8 plots out of these 18 plots and conducts a ground count on these 8 plots. Let x denote the number of dead trees in the plot by photo count and y the number of dead trees by ground count. The data are given as:

Plot	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
x'	5	7	10	6	7	9	3	6	8	11	5	9	12	13	3	20	15	4

Out of these 18 plots, 8 are randomly selected and a ground count is conducted.

Plot	2	3	5	6	12	15	16	17
x	7	10	7	9	9	3	20	15
y	9	13	10	11	10	4	25	17
$y-rx$	0.3375	0.6250	1.3375	-0.1375	-1.1375	0.2875	0.2500	-1.5625


3.2.1 Estimate the total number of dead trees in the 400-acre area. [6]

3.2.2 Compute the ratio estimate for the population total. [6]

3.2.3 Compute the estimated variance of the ratio estimator [8]

Question 4 [25 marks]

4.1 A mathematics achievement test was given to 486 students prior to entering a certain college who then took a calculus class. A simple random sampling of 10 students are selected and their calculus score recorded. It is known that the average achievement test score for the 486 students was 52. The scatterplot of the 10 samples are given on page 4 and the data follows.



The scatter plot shows that there is a strong positive linear relationship.

$$\hat{\mu}_x = \bar{y} + b(\mu_x - \bar{x}) = a + b\mu_x$$

Student	Achievement test score X	Calculus score Y
1	39	65
2	43	78
3	21	52
4	64	82
5	57	92
6	47	89
7	28	73
8	75	98
9	34	56
10	52	75

The regression equation is
 $Y = 40.8 + 0.766 X$

Predictor	Coef	StDev	T	P
Constant	40.784	8.507	4.79	0.001
X	0.7656	0.1750	4.38	0.002

S = 8.704 R-Sq = 70.5% R-Sq(adj) = 66.8%

Analysis of Variance					
Source	DF	SS	MS	F	P
Regression	1	1450.0	1450.0	19.14	0.002
Resid. Err	8	606.0	75.8		
Total	9	2056.0			

4.1.1 Using the results from the output above, calculate the regression estimate. [3]

4.1.2 What is the variance of the regression estimate? [5]


4.1.3 Calculate the approximate 95% Confidence Limits for μ . [7]

4.2 A population of 20000 farms were divided into 30 clusters. Sample 3000 farms from 10 clusters using Probability Proportional to size (PPS) by completing the table below in your answer sheet. [10]

A	B	C	D	E	F	G	H
Cluster	Size	Cumulative sum	Clusters sampled	Prob 1	Individuals per cluster	Prob 2	Overall weight
1	1028		905		300		
2	555						
3	390						
4	1309		2905		300		
5	698						
6	907						
7	432		4905		300		
8	897						
9	677						
10	501		6905		300		
11	867						
12	867		8905		300		
13	1002						
14	1094		10905		300		
15	668						
16	500						
17	835		12905		300		
18	396						
19	630						
20	483						
21	319		14905		300		
22	569						
23	987						
24	598		16905		300		
25	375						
26	387						
27	465						
28	751		18905		300		
29	365						
30	448	20000					

*****END OF EXAMINATION*****

Numbers in each row of the table are values on a t -distribution with (df) degrees of freedom for selected right-tail (greater-than) probabilities (p).



df/p	0.40	0.25	0.10	0.05	0.025	0.01	0.005	0.0005
1	0.324920	1.000000	3.077684	6.313752	12.70620	31.82052	63.65674	636.6192
2	0.288675	0.816497	1.885618	2.919986	4.30265	6.96456	9.92484	31.5991
3	0.276671	0.764892	1.637744	2.353363	3.18245	4.54070	5.84091	12.9240
4	0.270722	0.740697	1.533206	2.131847	2.77645	3.74695	4.60409	8.6103
5	0.267181	0.726687	1.475884	2.015048	2.57058	3.36493	4.03214	6.8688
6	0.264835	0.717558	1.439756	1.943180	2.44691	3.14267	3.70743	5.9588
7	0.263167	0.711142	1.414924	1.894579	2.36462	2.99795	3.49948	5.4079
8	0.261921	0.706387	1.396815	1.859548	2.30600	2.89646	3.35539	5.0413
9	0.260955	0.702722	1.383029	1.833113	2.26216	2.82144	3.24984	4.7809
10	0.260185	0.699812	1.372184	1.812461	2.22814	2.76377	3.16927	4.5869
11	0.259556	0.697445	1.363430	1.795885	2.20099	2.71808	3.10581	4.4370
12	0.259033	0.695483	1.356217	1.782288	2.17881	2.68100	3.05454	4.3178
13	0.258591	0.693829	1.350171	1.770933	2.16037	2.65031	3.01228	4.2208
14	0.258213	0.692417	1.345030	1.761310	2.14479	2.62449	2.97684	4.1405
15	0.257885	0.691197	1.340606	1.753050	2.13145	2.60248	2.94671	4.0728
16	0.257599	0.690132	1.336757	1.745884	2.11991	2.58349	2.92078	4.0150
17	0.257347	0.689195	1.333379	1.739607	2.10982	2.56693	2.89823	3.9651
18	0.257123	0.688364	1.330391	1.734064	2.10092	2.55238	2.87844	3.9216
19	0.256923	0.687621	1.327728	1.729133	2.09302	2.53948	2.86093	3.8834
20	0.256743	0.686954	1.325341	1.724718	2.08596	2.52798	2.84534	3.8495
21	0.256580	0.686352	1.323188	1.720743	2.07961	2.51765	2.83136	3.8193
22	0.256432	0.685805	1.321237	1.717144	2.07387	2.50832	2.81876	3.7921
23	0.256297	0.685306	1.319460	1.713872	2.06866	2.49987	2.80734	3.7676
24	0.256173	0.684850	1.317836	1.710882	2.06390	2.49216	2.79694	3.7454
25	0.256060	0.684430	1.316345	1.708141	2.05954	2.48511	2.78744	3.7251
26	0.255955	0.684043	1.314972	1.705618	2.05553	2.47863	2.77871	3.7066
27	0.255858	0.683685	1.313703	1.703288	2.05183	2.47266	2.77068	3.6896
28	0.255768	0.683353	1.312527	1.701131	2.04841	2.46714	2.76326	3.6739
29	0.255684	0.683044	1.311434	1.699127	2.04523	2.46202	2.75639	3.6594
30	0.255605	0.682756	1.310415	1.697261	2.04227	2.45726	2.75000	3.6460
z	0.253347	0.674490	1.281552	1.644854	1.95996	2.32635	2.57583	3.2905
CI	————	————	80%	90%	95%	98%	99%	99.9%

Standard Normal Probabilities




Table entry for z is the area under the standard normal curve to the left of z .

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

Standard Normal Probabilities




Table entry for z is the area under the standard normal curve to the left of z .

[illegible]