



**NAMIBIA UNIVERSITY
OF SCIENCE AND TECHNOLOGY**

**FACULTY OF HEALTH, NATURAL RESOURCES AND APPLIED SCIENCES
SCHOOL OF NATURAL AND APPLIED SCIENCES
DEPARTMENT OF MATHEMATICS, STATISTICS AND ACTUARIAL SCIENCE**

QUALIFICATION:	BACHELOR OF SCIENCE IN APPLIED MATHEMATICS AND STATISTICS		
QUALIFICATION CODE:	07BSAM	LEVEL:	7
COURSE CODE:	NUM701S	COURSE NAME:	NUMERICAL METHODS 1
SESSION:	JULY 2023	PAPER:	THEORY
DURATION:	3 HOURS	MARKS:	100

SUPPLEMENTARY /SECOND OPPORTUNITY EXAMINATION QUESTION PAPER	
EXAMINERS	Dr S. N. NEOSI NGUETCHUE AND G. S. MBOKOMA
MODERATOR:	Prof S. S. MOTSA

INSTRUCTIONS
<ol style="list-style-type: none">1. Answer ALL the questions in the booklet provided.2. Show clearly all the steps used in the calculations. All numerical results must be given using 4 decimals where necessary unless mentioned otherwise.3. All written work must be done in blue or black ink and sketches must be done in pencil.

PERMISSIBLE MATERIALS

1. Non-programmable calculator without a cover.

THIS QUESTION PAPER CONSISTS OF 3 PAGES (Including this front page)

Attachments

None

Problem 1 [29 marks]

1-1. Consider the equation $f(x) = e^x - x^2 + 16 \sin(x) - 5 = 0$.

1-1-1. Show that $f(x) = 0$ has a unique solution α in $[0, 1]$. [6]

1-1-2. Use the Bisection method with initial interval $[0, 1]$ to find the approximate solution to α at the third iteration. [8]

1-2. Write down Newton's algorithm to approximate the root of a continuous function h in $[a, b]$ after n iterations. [3]

1-3. Suppose that $g : [a, b] \rightarrow [a, b]$ is continuous on the real interval $[a, b]$ and is a contraction in the sense that there exists a constant $\lambda \in (0, 1)$ such that

$$|g(x) - g(y)| \leq \lambda|x - y|, \text{ for all } x, y \in [a, b].$$

Prove that there exists a unique fixed point in $[a, b]$ and that the fixed point iteration $x_{n+1} = g(x_n)$ converges to it for any $x_0 \in [a, b]$. Also, prove that the error is reduced by a factor of at least λ from each iteration to the next. [12]

Problem 2. [39 marks]

2-1. Write down in details the formulae of the Lagrange and Newton's form of the polynomial that interpolates the set of data points $(x_0, f(x_0)), (x_1, f(x_1)), \dots, (x_n, f(x_n))$. [7]

2-2. Use the results in 2-1. to determine the Lagrange and Newton's form of the polynomial that interpolates the set of data points $(1, 1), (2, 5)$ and $(3, 15)$. [18]

2-3. Determine the error term for the formula [14]

$$f'''(x) \approx \frac{1}{2h^3}[3f(x+h) - 10f(x) + 12f(x-h) - 6f(x-2h) + f(x-3h)]$$

Problem 3. [32 marks]

3-1. State the Improved-Euler's algorithm and indicate its order of accuracy. [4]

3-2. Write down the fourth-order Runge-Kutta (RK4) method's algorithm for the following specific problem after n steps [8]

$$y'(t) = y - t^2 + 1, \quad y(0) = 2$$

3-3. In the kingdom of Bana, king Happi The First asked one of his subjects, a prominent mathematician to solve the above IVP using the fourth-order Runge-Kutta (RK4) method. He displayed the results in the form of the following table and purposely skipped some entries.

k	t_k	k_1	k_2	k_3	k_4	y_k
1	0.08	3.0	3.11840		3.24345	2.24969
2	0.16		3.36502		3.49368	
3		3.49351	3.61885			2.80885
4		3.75125		3.88567	4.01730	
5	0.4		4.15061		4.29200	

Compute **only the missing values** by the means of the given ones (don't re-compute them!!).

[20]

TOTAL: 100 marks

God bless you !!!